

NUMERICAL SOLUTION OF A NONLINEAR REACTION-DIFFUSION PROBLEM IN THE CASE OF HS-REGIME

MARIE-NOELLE LE ROUX

ABSTRACT. In this paper, the author propose a numerical method to compute the solution of the Cauchy problem: $w_t - (w^m w_x)_x = w^p$, the initial condition is a nonnegative function with compact support, $m > 0$, $1 < p < m + 1$. The problem is split in two parts: A hyperbolic term solved by using the Hopf and Lax formula and a parabolic term solved by a backward linearized Euler method in time and a finite element method in space. Estimates of the numerical solution are obtained and it is proved that any numerical solution is unbounded.

1. INTRODUCTION

In this paper, we study a numerical method to compute the solution of the Cauchy problem :

$$(1.1) \quad \begin{aligned} w_t - (w^m w_x)_x &= w^p, \quad t > 0, \quad x \in \mathbb{R} \\ w(x, 0) &= w_0(x) \geq 0, \quad x \in \mathbb{R} \end{aligned}$$

w_0 is a function with compact support, $m > 0$, $1 < p < m + 1$.

Samarskii et al [10], see also [1], [2], [3], [6], [5], [4], [9] have obtained theoretical results on this problem. In the case $1 < p < m + 1$, the numerical solution blows up in finite time and there is no localization (HS-regime) that is $u(t, x) \rightarrow \infty$ in $\mathbb{R}$ if $t \rightarrow T_0$.

A numerical method to solve (1.1) has been proposed in the case of S-regime ($p = m + 1$) in [7] and in the case of LS-regime ($p > m + 1$) [8]. If we denote $\omega_L = \{x \in \mathbb{R} / u(T_0^-, x) = \infty\}$, in the case of S-regime, $L^* = \text{mes}(\omega_L)$ is positive while in the case of LS-regime, $L^* = 0$. The problem is solved by using a splitting method; for that, it is more convenient to work with the function $u = w^m$. Problem (1.1) may be written:

$$(1.2) \quad \begin{aligned} u_t - \frac{1}{m} u_x^2 - u u_{xx} &= m u^{q+1}, \quad t > 0, \quad x \in \mathbb{R} \\ u(x, 0) &= u_0(x) = w_0^m(x), \quad x \in \mathbb{R} \end{aligned}$$

with $q = \frac{p-1}{m}$, $m > 0$, $q < 1$.

This problem is split in two parts: a hyperbolic problem which will be solved exactly at the nodes at each time step and allows the extension of the domain and a parabolic problem which will be solved by a backward linearized Euler method which allows the blow up of the solution.

In [7], [8], the convergence of the scheme has been proved in the cases $q = 1$ and $q > 1$. It has also been proved that for $q = 1$, the numerical solution blows up in finite time for any initial condition

and that its support remains bounded if the initial condition is smaller than a self-similar solution. In the case, $1 \leq q < \frac{m+2}{m}$, any numerical solution is unbounded while for $q > \frac{m+2}{m}$ ($p > m + 3$) if the initial condition is sufficiently small, a global solution exists and if $q \geq \frac{m+2}{m}$ for large initial condition, the solution blows up in a finite time. We observe numerically that in any case, the unbounded solution is strictly localized and blows up in one point and that for $q = \frac{m+2}{m}$ also, if the initial condition is sufficiently small, a global solution exists.

Here, we generalize this method to the case $q < 1$.

An outline of the paper is as follows:

In Section 2, we present the numerical scheme. In Section 3, we obtain estimates of the approximate solution and of its derivative in space. In Section 4, we prove that if $q < 1$, ($p < m + 1$) any numerical solution is unbounded

2. DEFINITION OF THE NUMERICAL SOLUTION

In order to solve problem (1.2), we separate it in two parts: a hyperbolic problem

$$(2.1) \quad u_t - \frac{1}{m}(u_x)^2 = 0, \quad x \in \mathbb{R}, \quad t > 0$$

and a parabolic problem

$$(2.2) \quad u_t - uu_{xx} = mu^{q+1}, \quad x \in \mathbb{R}, \quad t > 0$$

We denote by Δt_n the time increment between the time levels t_n and t_{n+1} , $n \geq 0$ and by u_h^n the approximate solution at the time level t_n . This solution will be in a finite-dimensional space which will be defined below.

Without loss of generality, we can assume that the initial condition is a continuous function with a symmetric compact support $[-s_0, s_0]$. Let $N \in \mathbb{N}$, the space step h is defined by $h = \frac{s_0}{N}$; we note $x_i = ih$, $i \in \mathbb{Z}$, $I_i = (x_{i-1}, x_i)$ and we define the finite dimensional space V_h^0 by

$$(2.3) \quad V_h^0 = \left\{ \phi_h \in C^0(\mathbb{R}) / \phi_h(x) = 0, \quad x \notin [-s_0, +s_0], \quad \phi_h|_{I_i} \in P_1, \quad i = -N + 1, N \right\}$$

For $\phi_h \in V_h^0$, we note $\phi_i = \phi_h(x_i)$, $i \in \mathbb{Z}$ and for any function $v \in C^0(\mathbb{R})$ with compact support $[-s_0, +s_0]$, we define its interpolate by $\pi_h^0 v \in V_h^0$ and $\pi_h^0 v(x_i) = v(x_i)$, $i \in \mathbb{Z}$.

The support of the solution u_h^n will be denoted $[-s_-^n, s_+^n]$ and will be computed at each time level by solving (2.1).

We denote

$$N_-^n = \left\lceil \frac{s_-^n}{h} \right\rceil, \quad N_+^n = \left\lceil \frac{s_+^n}{h} \right\rceil, \quad h_-^n = s_-^n - (N_-^n - 1)h, \quad h_+^n = s_+^n - (N_+^n - 1)h$$

(for $x \in \mathbb{R}$, $[x]$ denotes the greatest integer less than x), so we get: $h \leq h_-^n, h_+^n < 2h$.

We then define the finite-dimensional space V_h^n by:

$$V_h^n = \left\{ \begin{array}{l} \phi_h \in C^0(\mathbb{R}) / \phi_h(x) = 0, \ x \notin [-s_-^n, s_+^n], \\ \phi_h|_{I_i} \in P_1, i = -N_-^n + 2, N_+^n - 1 \\ \phi_h|_{(-s_-^n, x_{-N_-^n+1})}, \phi_h|_{(x_{N_+^n-1}, s_+^n)} \in P_1 \end{array} \right\}$$

and denote by $\pi_h^n v$ the Lagrange interpolate in V_h^n of a function $v \in C^0(\mathbb{R})$ with a compact support in $[-s_-^n, s_+^n]$.

The solution u_h^{n+1} at the time level t_{n+1} is computed in two steps: knowing u_h^n , we compute first an approximate solution of (2.1) which will be denoted $u_h^{n+\frac{1}{2}}$; then starting with this intermediate value, we compute an approximate solution of (2.2) at time level t_{n+1} .

Then if S is the semi-group operator associated with (2.1), we define

$$u^{n+\frac{1}{2}} = S(\Delta t_n) u_h^n.$$

The support of this function will be denoted $[-s_-^{n+1}, s_+^{n+1}]$ and the interpolate of $u^{n+\frac{1}{2}}$ in V_h^{n+1} will be denoted $u_h^{n+\frac{1}{2}} = \pi_h^{n+1} u^{n+\frac{1}{2}}$.

Then starting with $u_h^{n+\frac{1}{2}}$, the function u_h^{n+1} is obtained by solving (2.2) with a backward linearized Euler method in time and a P_1 -finite element method with numerical integration in space. This function has the same support as $u_h^{n+\frac{1}{2}}$.

2.1. Computation of the solution of the hyperbolic problem.

The hyperbolic problem is independent of q ; we use the same method as in [7] for the case $q = 1$. It is not necessary in this case to use P_2 -interpolation on the last interval since there no localization. We use the Hopf and Lax formula which gives explicitly the solution to (2.1) with the starting data u_h^n at the time level $t = t_n$. Here, we simply recall the results obtained in [7]

We define the piecewise constant function v_h^n by

$$v_i^n = \frac{u_i^n - u_{i-1}^n}{h} \text{ on } I_i, \ -N_-^n + 2 \leq i \leq N_+^n - 1$$

$$v_{N_+^n}^n = -\frac{u_{N_+^n-1}^n}{h_+^n} \text{ on } (x_{N_+^n-1}, s_+^n)$$

$$v_{-N_-^n+1}^n = \frac{u_{-N_-^n+1}^n}{h_-^n} \text{ on } (-s_-^n, x_{-N_-^n+1}).$$

Let us denote $r_n = \frac{\Delta t_n}{h}$, $\|v\|_s = \|v\|_{L^s(\mathbb{R})}$, $s > 0$.

Proposition 2.1. *If the following stability condition*

$$(2.4) \quad r_n \|v_h^n\|_\infty \leq \frac{m}{2}$$

is satisfied, then the solution u_h^{n+1} of (2.1) is defined by

$$u_i^{n+\frac{1}{2}} = u_i^n + \frac{\Delta t_n}{m} \left(\max(0, -v_i^n, v_{i+1}^n) \right)^2, \quad -N_-^n + 1 \leq i \leq N_+^n - 1$$

$$s_+^{n+1} = s_+^n - \frac{\Delta t_n}{m} v_{N_+^n}^n$$

$$s_-^{n+1} = s_-^n + \frac{\Delta t_n}{m} v_{-N_-^n+1}^n$$

If $N_+^{n+1} = N_+^n + 1$, we get

$$u_{N_+^n}^{n+\frac{1}{2}} = \left(1 - \frac{h}{h_+^n} \right) u_{N_+^n-1}^n + \frac{\Delta t_n}{m} \left(v_{N_+^n}^n \right)^2$$

and we get analogous formula at the other end of the support.

2.2. Computation of the parabolic problem.

The approximate solution at t_{n+1} is now obtained by solving problem (2.2).

We introduce the approximate scalar product on V_h^{n+1} : $\forall \phi_h, \psi_h \in V_h^{n+1}$,

$$(\phi_h, \psi_h)_h = \frac{1}{2}(h_-^{n+1} + h)\phi_{-N_-^{n+1}+1}\psi_{-N_-^{n+1}+1} + h \sum_{i=-N_-^{n+1}+2}^{i=N_+^{n+1}-2} \phi_i \psi_i + \frac{1}{2}(h_+^{n+1} + h)\phi_{N_+^{n+1}-1}\psi_{N_+^{n+1}-1}.$$

We define u_h^{n+1} as the solution of the following problem:

$$(2.5) \quad \forall \phi_h \in V_h^{n+1}, \begin{cases} (u_h^{n+1}, \phi_h)_h + \Delta t_n ((u_h^{n+1})_x, (u_h^{n+\frac{1}{2}} \phi_h)_x) = \\ (u_h^{n+\frac{1}{2}}, \phi_h)_h + m q \Delta t_n \left(\pi_h^{n+1} \left(\left(u_h^{n+\frac{1}{2}} \right)^q u_h^{n+1} \right), \phi_h \right)_h \\ + m(1-q) \Delta t_n \left(\pi_h^{n+\frac{1}{2}} \left(\left(u_h^{n+\frac{1}{2}} \right)^{q+1} \right), \phi_h \right)_h \end{cases}$$

The second member of (2.2) is splitted in two parts in order to obtain the L^∞ -estimate of u_h^{n+1} . This equation may be written:

$$\left(1 - m q \Delta t_n \left(u_i^{n+\frac{1}{2}} \right)^q \right) u_i^{n+1} + \frac{\Delta t_n}{h^2} u_i^{n+\frac{1}{2}} (2u_i^{n+1} - u_{i-1}^{n+1} - u_{i+1}^{n+1}) =$$

$$u_i^{n+\frac{1}{2}} + m(1-q)\Delta t_n \left(u_i^{n+\frac{1}{2}}\right)^{q+1}, \quad -N_-^{n+1} + 2 \leq i \leq N_+^{n+1} - 2$$

$$\left(1 - mq\Delta t_n \left(u_{N_+^{n+1}-1}^{n+\frac{1}{2}}\right)^q\right) u_{N_+^{n+1}-1}^{n+1} +$$

$$\frac{\Delta t_n}{h} u_{N_+^{n+1}-1}^{n+\frac{1}{2}} \left(\frac{2}{h_+^{n+1}} u_{N_+^{n+1}-1}^{n+1} - \frac{2}{h_+ + h_+^{n+1}} u_{N_+^{n+1}-2}^{n+1} \right)$$

$$= u_{N_+^{n+1}-1}^{n+\frac{1}{2}} + m(1-q)\Delta t_n \left(u_{N_+^{n+1}-1}^{n+\frac{1}{2}}\right)^{q+1}$$

$$\left(1 - mq\Delta t_n \left(u_{-N_-^{n+1}+1}^{n+\frac{1}{2}}\right)^q\right) u_{-N_-^{n+1}+1}^{n+1} + \frac{\Delta t_n}{h} u_{-N_-^{n+1}+1}^{n+\frac{1}{2}} \left(\frac{2}{h_-^{n+1}} u_{-N_-^{n+1}+1}^{n+1} - \frac{2}{h_-^{n+1} + h} u_{-N_-^{n+1}+2}^{n+1} \right)$$

$$= u_{-N_-^{n+1}+1}^{n+\frac{1}{2}} + m(1-q)\Delta t_n \left(u_{-N_-^{n+1}+1}^{n+\frac{1}{2}}\right)^{q+1}$$

We get immediately the result:

Proposition 2.2. *If the hypotheses of proposition 2.2 are satisfied and if $u_h^{n+\frac{1}{2}}$ satisfies*

$$(2.6) \quad mq\Delta t_n \left\| u_h^{n+\frac{1}{2}} \right\|_\infty^q < 1$$

then the solution u_h^{n+1} of (2.2) is unique and nonnegative.

2.3. Properties of the scheme.

Lemma 2.3. *If the hypotheses of proposition 2.2 are satisfied, then the following estimate holds:*

$$(2.7) \quad \|u_h^{n+1}\|_\infty \leq \frac{\|u_h^n\|_\infty (1 + m(1-q)\Delta t_n \|u_h^n\|_\infty^q)}{1 - mq\Delta t_n \|u_h^n\|_\infty^q}$$

Proof: We get immediately from the Hopf and Lax formula that $\left\|u_h^{n+\frac{1}{2}}\right\|_\infty \leq \|u_h^n\|_\infty$. If we denote i_0 the index such that $u_{i_0}^{n+1} = \|u_h^{n+1}\|_\infty$, we get from (2.2), (2.2), (2.2) that

$$u_{i_0}^{n+1} \leq u_{i_0}^{n+\frac{1}{2}} \frac{1 + m(1-q)\Delta t_n \left(u_{i_0}^{n+\frac{1}{2}}\right)^q}{1 - mq\Delta t_n \|u_h^n\|_\infty^q}$$

which proves the lemma.

We deduce the following theorem:

Theorem 2.4. *Under the hypotheses of proposition 2.2, the numerical solution exists at least until the time*

$$(2.8) \quad T_1 = \frac{1}{mq \|u_h^0\|_\infty^q}$$

and the following estimate holds:

$$(2.9) \quad \|u_h^n\|_\infty \leq \frac{\|u_h^0\|_\infty}{(1 - mqt_n \|u_h^0\|_\infty^q)^{\frac{1}{q}}}$$

Proof: This result is proved recurrently. It is true for $n = 0$. If we suppose that we have estimate (2.9) at the time level t_n , we get from (2.7), at the time t_{n+1} :

$$\|u_h^{n+1}\|_\infty \leq \|u_h^n\|_\infty \frac{1 + m(1-q)\Delta t_n \|u_h^n\|_\infty^q}{1 - mq\Delta t_n \|u_h^n\|_\infty^q}$$

or

$$\|u_h^{n+1}\|_\infty \leq \|u_h^0\|_\infty \frac{1 - mqt_n \|u_h^0\|_\infty^q + m(1-q)\Delta t_n \|u_h^0\|_\infty^q}{(1 - mqt_{n+1} \|u_h^0\|_\infty^q) (1 - mqt_n \|u_h^0\|_\infty^q)^{\frac{1}{q}}}$$

The inequality (2.9) will be satisfied at the time t_{n+1} if :

$$\begin{aligned} & (1 - mqt_{n+1} \|u_h^0\|_\infty^q + m\Delta t_n \|u_h^0\|_\infty^q) \\ & \leq (1 - mqt_n \|u_h^0\|_\infty^q)^{\frac{1}{q}} (1 - mqt_{n+1} \|u_h^0\|_\infty^q)^{1-\frac{1}{q}} \end{aligned}$$

By using the Taylor formula, we get:

$$(1 - mqt_n \|u_h^0\|_\infty^q)^{\frac{1}{q}} - (1 - mqt_{n+1} \|u_h^0\|_\infty^q)^{\frac{1}{q}} \geq m\Delta t_n \|u_h^0\|_\infty^q (1 - mqt_{n+1} \|u_h^0\|_\infty^q)^{\frac{1}{q}-1}$$

and the inequality (2.3) is satisfied:

Lemma 2.5. *Under the hypothesis of Proposition (2.2), we have the estimate:*

$$\|v_h^n\|_\infty \leq C$$

for $t_n \leq T < T_1$ where C is a constant depending on T and u_0 .

Proof: We have the inequality([7]): $\left\|v_h^{n+\frac{1}{2}}\right\|_\infty \leq \|v_h^n\|_\infty$.

It remains to estimate $\left\|v_h^{n+1}\right\|_\infty$.

From (2.2), we deduce the following equation satisfied by v_h^{n+1} :

$$\begin{aligned} & \left(1 - mq\Delta t_n \left(u_i^{n+\frac{1}{2}}\right)^q\right) v_i^{n+1} + \frac{\Delta t_n}{h^2} \left(v_i^{n+1} \left(u_i^{n+\frac{1}{2}} + u_{i-1}^{n+\frac{1}{2}}\right) - v_{i-1}^{n+1} u_{i-1}^{n+\frac{1}{2}} - v_{i+1}^{n+1} u_i^{n+\frac{1}{2}}\right) \\ &= v_i^{n+\frac{1}{2}} + mq \frac{\Delta t_n}{h} \left(\left(u_i^{n+\frac{1}{2}}\right)^q - \left(u_{i-1}^{n+\frac{1}{2}}\right)^q\right) u_{i-1}^{n+1} + m(1-q) \frac{\Delta t_n}{h} \left(\left(u_i^{n+\frac{1}{2}}\right)^{q+1} - \left(u_{i-1}^{n+\frac{1}{2}}\right)^{q+1}\right) \end{aligned}$$

$$-N_-^{n+1} + 3 \leq i \leq N_+^{n+1} - 2$$

and we have analogous inequalities for $i = N_+^{n+1} - 1, N_+^{n+1}, -N_-^{n+1} + 2, -N_-^{n+1} + 1$.

By using (2.2) for $i - 1$, we can replace u_{i-1}^{n+1} in the second member by its expression in function of the values of $u_h^{n+\frac{1}{2}}$ and v_h^{n+1} and we get:

$$\begin{aligned} & \left(1 - mq\Delta t_n \left(u_i^{n+\frac{1}{2}}\right)^q\right) v_i^{n+1} + \\ & \frac{\Delta t_n}{h^2} \left((v_i^{n+1} - v_{i+1}^{n+1}) u_i^{n+\frac{1}{2}} + (v_i^{n+1} - v_{i-1}^{n+1}) u_i^{n+\frac{1}{2}} \frac{1 - mq\Delta t_n \left(\left(u_i^{n+\frac{1}{2}}\right)^q - \left(u_{i-1}^{n+\frac{1}{2}}\right)^q\right)}{1 - mq\Delta t_n \left(u_i^{n+\frac{1}{2}}\right)^q} \right) \\ &= v_i^{n+\frac{1}{2}} + mq \frac{\Delta t_n}{h} \left(\left(u_i^{n+\frac{1}{2}}\right)^q - \left(u_{i-1}^{n+\frac{1}{2}}\right)^q\right) u_i^{n+\frac{1}{2}} \frac{1 + m(1-q)\Delta t_n \left(u_{i-1}^{n+\frac{1}{2}}\right)^q}{1 - mq\Delta t_n \left(u_{i-1}^{n+\frac{1}{2}}\right)^q} \\ & \quad + m(1-q) \frac{\Delta t_n}{h} \left(\left(u_i^{n+\frac{1}{2}}\right)^{q+1} - \left(u_{i-1}^{n+\frac{1}{2}}\right)^{q+1}\right) \end{aligned}$$

Let i_0 the index such that $|v_{i_0}^{n+1}| = \|v_h^{n+1}\|_\infty$. From the previous equality, we get:

$$\begin{aligned} (1 - mq\Delta t_n \|u_h^n\|_\infty^q) |v_{i_0}^{n+1}| &\leq |v_{i_0}^{n+\frac{1}{2}}| + m(1-q) \frac{\Delta t_n}{h} \left| \left(u_{i_0}^{n+\frac{1}{2}}\right)^{q+1} - \left(u_{i_0-1}^{n+\frac{1}{2}}\right)^{q+1} \right| \\ &\quad + mq \frac{\Delta t_n}{h} \left| \left(u_{i_0}^{n+\frac{1}{2}}\right)^q - \left(u_{i_0-1}^{n+\frac{1}{2}}\right)^q \right| u_{i_0}^{n+\frac{1}{2}} \frac{1 + m(1-q)\Delta t_n \|u_h^{n+1}\|_\infty}{1 - mq\Delta t_n \|u_h^n\|_\infty^q} \end{aligned}$$

We easily obtain:

$$\frac{1}{h} \left| \left(u_{i_0}^{n+\frac{1}{2}}\right)^{q+1} - \left(u_{i_0-1}^{n+\frac{1}{2}}\right)^{q+1} \right| \leq (q+1) \|u_h^n\|_\infty^q |v_{i_0}^{n+\frac{1}{2}}|$$

and

$$\frac{1}{h} \left| \left(u_{i_0}^{n+\frac{1}{2}}\right)^q - \left(u_{i_0-1}^{n+\frac{1}{2}}\right)^q \right| u_{i_0}^{n+\frac{1}{2}} \leq |v_{i_0}^{n+\frac{1}{2}}| \|u_h^n\|_\infty^q$$

and we get:

$$\begin{aligned} (1 - mq\Delta t_n \|u_h^n\|_\infty^q) \|v_h^{n+1}\|_\infty &\leq \\ \|v_h^n\|_\infty &\left(1 + m(1-q^2)\Delta t_n \|u_h^n\|_\infty^q + mq\Delta t_n \|u_h^n\|_\infty^q \frac{1 + m(1-q)\Delta t_n \|u_h^{n+1}\|_\infty^q}{1 - mq\Delta t_n \|u_h^n\|_\infty^q} \right) \end{aligned}$$

By (2.9), we get easily there exist a positive constant C depending on $m, q, T, \|u_h^0\|_\infty$ such that

$$\|v_h^{n+1}\|_\infty \leq (1 + C\Delta t_n) \|v_h^n\|_\infty$$

Hence for $t_n \leq T < T_1$, we get $\|v_h^n\|_\infty \leq C$.

Lemma 2.6. *Under the hypotheses of Proposition (2.2), we have the estimate:*

$$(2.10) \quad \|v_h^n\|_1 \leq C$$

for $t_n < T < T_1$ where C is a constant depending on T and u_0 .

Proof: From the properties of the semigroup operator S [7], we get: $\|v_h^{n+\frac{1}{2}}\|_1 \leq \|v_h^n\|_1$, and by using the equations satisfied by $v_i^{n+\frac{1}{2}}$, $-N_-^{n+1} \leq i \leq N_+^{n+1}$, we obtain:

$$\left(1 - mq\Delta t_n \left\|u_h^{n+\frac{1}{2}}\right\|_\infty^q\right) \|v_h^{n+1}\|_1 \leq \left\|v_h^{n+\frac{1}{2}}\right\|_1 \left(1 + m\Delta t_n \left\|u_h^{n+\frac{1}{2}}\right\|_\infty^q \left(q^2 \|u_h^{n+1}\|_\infty + (1 - q^2) \left\|u_h^{n+\frac{1}{2}}\right\|_\infty\right)\right)$$

and we immediately deduce the estimate (2.10).

In this case, since $q < 1$, the variation of v_h^n is not bounded.

3. BLOW-UP OF THE SOLUTION

In this part, we prove that for $q < 1$ the solution blows up in finite time.

3.1. Construction of unbounded solutions.

Define the function

$$\theta(x) = \begin{cases} 1 - \frac{x^2}{a^2}, & |x| \leq a \\ 0, & |x| \geq a \end{cases}$$

We note $\theta_h = \pi_h \theta$. If the initial condition is $u_h^0 = \frac{\lambda}{T^{\frac{1}{q}}} \theta_h(\xi^0)$ with $\xi^0 = \frac{x}{T^{\frac{q-1}{2q}}}$, we prove that it is possible to choose λ and a in such a manner that $u_h^n(x) \geq \frac{\lambda}{(T-t_n)^{\frac{1}{q}}} \theta_h(\xi_n)$, with $\xi_n = x(T-t_n)^{\frac{1-q}{2q}}$ and then the numerical solution blows up in finite time.

We denote $\zeta_n = (T-t_n)^{\frac{q-1}{2q}}$, $\hat{u}_h^n(x) = \frac{\lambda}{(T-t_n)^{\frac{1}{q}}} \theta_h(\xi_n)$.

The support of $\hat{u}_h^n$ is $] -a\zeta_n, a\zeta_n[$; its length is increasing with the time.

If $\hat{u}_h^n \leq u_h^n$, we get $\hat{u}_h^{n+\frac{1}{2}} \leq u_h^{n+\frac{1}{2}}$.

The support of $\hat{u}_h^{n+\frac{1}{2}}$ is $[-\hat{s}_-^{n+1}, \hat{s}_+^{n+1}]$ with $\hat{s}_+^{n+1} \leq s^{n+1}$, $\hat{s}_-^{n+1} \leq s_-^{n+1}$.

Since $\hat{u}_h^n$ is a symmetric function, it is sufficient to study the case $x \geq 0, (i \geq 0)$.

We get for $i \geq 0$,

$$\hat{u}_i^{n+\frac{1}{2}} = \hat{u}_i^n + \frac{\Delta t_n}{m} (\hat{v}_i^n)^2$$

for i such that $x_i \leq a\zeta_n$ since the function $\hat{u}_h^n$ is decreasing for $x \geq 0$ and we have:

$$\hat{v}_i^n = \frac{\hat{u}_i^n - \hat{u}_{i-1}^n}{h} = \frac{\lambda}{(T-t_n)^{\frac{1}{q}}} \frac{\theta_i^n - \theta_{i-1}^n}{h}$$

with $\theta_i^n = \theta(\xi_i^n)$.

Hence, we obtain:

$$\hat{u}_i^{n+\frac{1}{2}} = \frac{\lambda}{(T-t_n)^{\frac{1}{q}}} \left(\theta_i^n + \frac{4\lambda}{ma^2} \frac{\Delta t_n}{(T-t_n)^{\frac{1}{q}}} \left(1 - \theta_{i-\frac{1}{2}}^n \right) \right)$$

with $\theta_{i-\frac{1}{2}}^n = \theta(\xi_{i-\frac{1}{2}}^n)$, $\xi_{i-\frac{1}{2}}^n = \frac{1}{2} (\xi_i^n + \xi_{i-1}^n)$.

Then $\hat{u}_h^{n+1}$ will be a subsolution of (2.2) if

$$\begin{aligned} & \frac{\lambda}{(T-t_{n+1})^{\frac{1}{q}}} \left(1 - mq\Delta t_n \left(\hat{u}_i^{n+\frac{1}{2}} \right)^q \right) \theta_i^{n+1} + \frac{\lambda}{(T-t_{n+1})^{\frac{1}{q}}} \frac{\Delta t_n}{h^2} \hat{u}_i^{n+\frac{1}{2}} (2\theta_i^{n+1} - \theta_{i-1}^{n+1} - \theta_{i+1}^{n+1}) \\ & \leq \hat{u}_i^{n+\frac{1}{2}} + m(1-q)\Delta t_n \left(\hat{u}_i^{n+\frac{1}{2}} \right)^{q+1} \end{aligned}$$

By using the equality : $\frac{1}{h^2} (2\theta_i^{n+1} - \theta_{i-1}^{n+1} - \theta_{i+1}^{n+1}) = \frac{2}{a^2 \zeta_{n+1}^2}$

this inequality reduces after simplifications to

$$\begin{aligned} & \frac{\lambda}{(T-t_{n+1})^{\frac{1}{q}}} \theta_i^{n+1} \leq \hat{u}_i^{n+\frac{1}{2}} \left(1 - \frac{2\lambda\Delta t_n}{a^2(T-t_{n+1})} \right) + \frac{\lambda}{(T-t_{n+1})^{\frac{1}{q}}} mq\Delta t_n \left(\hat{u}_i^{n+\frac{1}{2}} \right)^q \theta_i^{n+1} \\ & + m(1-q)\Delta t_n \left(\hat{u}_i^{n+\frac{1}{2}} \right)^{q+1} \end{aligned}$$

Noting that $\theta_{i-\frac{1}{2}}^n = \theta_i^n + \frac{h}{a^2 \zeta_n} \xi_{i-\frac{1}{2}}^n = \theta_i^n + \eta_i^n$ with $|\eta_i^n| \leq \frac{h}{a\zeta_n}$, we get:

$$\hat{u}_i^{n+\frac{1}{2}} = \frac{\lambda}{(T-t_n)^{\frac{1}{q}}} \left(\theta_i^n \left(1 - \frac{4\lambda}{ma^2} \frac{\Delta t_n}{T-t_n} \right) + \frac{4\lambda}{ma^2} \frac{\Delta t_n}{T-t_n} (1 - \eta_i^n) \right)$$

and the inequality (3.1) becomes:

$$\begin{aligned} & \frac{\theta_i^{n+1}}{(T-t_{n+1})^{\frac{1}{q}}} \leq \frac{\theta_i^n}{(T-t_n)^{\frac{1}{q}}} \left(1 - \frac{4\lambda}{ma^2} \frac{\Delta t_n}{T-t_n} - \frac{2\lambda}{ma^2} \frac{\Delta t_n}{T-t_{n+1}} \left(1 - \frac{4\lambda}{ma^2} \frac{\Delta t_n}{T-t_n} \right) \right) \\ & + \frac{4\lambda}{ma^2} \frac{\Delta t_n}{(T-t_{n+1})^{\frac{q+1}{q}}} \left(1 - \frac{2\lambda}{ma^2} \frac{\Delta t_n}{T-t_{n+1}} \right) (1 - \eta_i^n) + mq \frac{\Delta t_n}{(T-t_{n+1})^{\frac{1}{q}}} \left(\hat{u}_i^{n+\frac{1}{2}} \right)^q \theta_i^{n+1} \\ & + m(1-q) \frac{\Delta t_n}{\lambda} \left(\hat{u}_i^{n+\frac{1}{2}} \right)^{q+1} \end{aligned}$$

By using the equality $\theta_i^{n+1} = \theta_i^n \frac{\zeta_n^2}{\zeta_{n+1}^2} + 1 - \frac{\zeta_n^2}{\zeta_{n+1}^2}$, this inequality reduces to:

$$\theta_i^n \left(1 + \frac{4\lambda}{ma^2} \frac{T-t_{n+1}}{T-t_n} + \frac{2\lambda}{a^2} \left(1 - \frac{4\lambda}{ma^2} \frac{\Delta t_n}{T-t_n} \right) \right)$$

$$\begin{aligned} &\leq \frac{4\lambda}{ma^2} \frac{T-t_{n+1}}{T-t_n} \left(1 - \frac{2\lambda}{a^2} \frac{\Delta t_n}{T-t_{n+1}} (1-\eta_i^n)\right) - \left(\frac{\zeta_{n+1}^2}{\zeta_n^2} - 1\right) \frac{T-t_n}{\Delta t_n} \\ &\quad + mq (T-t_{n+1})^{1-\frac{1}{q}} (T-t_n)^{\frac{1}{q}} \left(\hat{u}_i^{n+\frac{1}{2}}\right)^q \theta_i^{n+1} + m(1-q) \frac{T-t_{n+1}}{\lambda} (T-t_n)^{\frac{1}{q}} \left(\hat{u}_i^{n+\frac{1}{2}}\right)^{q+1}. \end{aligned}$$

If we denote $\mu_n = \frac{4\lambda}{ma^2} \frac{\Delta t_n}{T-t_n}$ since $\hat{u}_i^{n+\frac{1}{2}} \geq \frac{\lambda}{(T-t_n)^{\frac{1}{q}}} \theta_i^n (1-\mu_n)$, the preceding inequality will be satisfied if:

$$\begin{aligned} (3.1) \quad &\theta_i^n \left(1 + \frac{4\lambda}{ma^2} + \frac{2\lambda}{a^2} - \mu_n \left(1 + \frac{2\lambda}{a^2}\right)\right) \leq \left(\frac{4\lambda}{ma^2} - \mu_n \left(1 + \frac{2\lambda}{a^2}\right)\right) (1-\eta_i^n) - \left(\frac{\zeta_{n+1}^2}{\zeta_n^2} - 1\right) \frac{T-t_n}{\Delta t_n} \\ &\quad + mq \lambda^q \frac{\zeta_{n+1}^2}{\zeta_n^2} (\theta_i^n)^q (1-\mu_n)^q \left(\theta_i^n \frac{\zeta_n^2}{\zeta_{n+1}^2} + 1 - \frac{\zeta_n^2}{\zeta_{n+1}^2}\right) \\ &\quad + m(1-q) \lambda^q \frac{T-t_{n+1}}{T-t_n} (\theta_i^n)^{q+1} (1-\mu_n)^{q+1} \end{aligned}$$

From the stability condition, we get:

$$\mu_n \leq \frac{h}{a\zeta_n} \leq \frac{h}{a\zeta_0} = \delta$$

hence for h sufficiently small, we get: $\frac{4\lambda}{ma^2} - \mu_n \left(1 + \frac{2\lambda}{a^2}\right) > 0$, and $|\eta_i^n| \leq \delta$

So the inequality (3.1) will be satisfied if :

$$\begin{aligned} &\theta_i^n \left(1 + \frac{4\lambda}{ma^2} + \frac{2\lambda}{a^2} - \mu_n \left(1 + \frac{2\lambda}{a^2}\right)\right) \leq \left(\frac{4\lambda}{ma^2} - \mu_n \left(1 + \frac{2\lambda}{a^2}\right)\right) (1-\delta) - \left(\frac{\zeta_{n+1}^2}{\zeta_n^2} - 1\right) \frac{T-t_n}{\Delta t_n} \\ &\quad + mq \lambda^q (1-\mu_n)^q (\theta_i^n)^q \left(\theta_i^n + \frac{\zeta_{n+1}^2}{\zeta_n^2} - 1\right) + m(1-q) \lambda^q (1-\mu_n)^{q+1} (\theta_i^n)^{q+1} \frac{T-t_{n+1}}{T-t_n} \end{aligned}$$

Since $\theta_i^n \in (0, 1)$, we introduce the function $\Phi_n(y)$ defined on $(0, 1)$ by :

$$\begin{aligned} \Phi_n(y) &= m \lambda^q (1-\mu_n)^q y^{q+1} \left(q + (1-q)(1-\mu_n) \frac{T-t_{n+1}}{T-t_n}\right) \\ &\quad + \left(\frac{4\lambda}{ma^2} - \mu_n \left(1 + \frac{2\lambda}{a^2}\right)\right) (1-\delta) - \left(\frac{\zeta_{n+1}^2}{\zeta_n^2} - 1\right) \frac{T-t_n}{\Delta t_n} \end{aligned}$$

$$-y \left(1 + \frac{4\lambda}{ma^2} + \frac{2\lambda}{a^2} - \mu_n \left(1 + \frac{2\lambda}{a^2} \right) \right)$$

or $\Phi_n(y) = A\lambda^q y^{q+1} + C - By$
with

$$A = m(1 - \mu_n)^q \left(q + (1 - q)(1 - \mu_n) \frac{T - t_{n+1}}{T - t_n} \right)$$

$$B = 1 + \frac{4\lambda}{ma^2} + \frac{2\lambda}{a^2} - \mu_n \left(1 + \frac{2\lambda}{a^2} \right)$$

$$C = \left(\frac{4\lambda}{ma^2} - \mu_n \left(1 + \frac{2\lambda}{a^2} \right) \right) (1 - \delta) - \left(\frac{\zeta_{n+1}^2}{\zeta_n^2} - 1 \right) \frac{T - t_n}{\Delta t_n}$$

A sufficient condition to satisfy (3.1) is: $\Phi_n(y) \geq 0$, $y \in (0, 1)$.

We have $\Phi_n(0) = C$,

hence we get

$$\Phi_n(0) \geq \left(\frac{4\lambda}{ma^2} - \delta \left(1 + \frac{2\lambda}{a^2} \right) \right) (1 - \delta) - \left(\frac{\zeta_{n+1}^2}{\zeta_n^2} - 1 \right) \frac{T - t_n}{\Delta t_n}$$

But since $0 < q < 1$, we get:

$$0 \leq \frac{\zeta_{n+1}^2}{\zeta_n^2} - 1 \leq \frac{1 - q}{q} \frac{\Delta t_n}{T - t_{n+1}} \frac{\zeta_n^2}{\zeta_{n+1}^2}$$

and

$$C \geq \left(\frac{4\lambda}{ma^2} - \delta \left(1 + \frac{2\lambda}{a^2} \right) \right) (1 - \delta) - \frac{1 - q}{q}.$$

So, if the quantity $\frac{\lambda}{a^2}$ is such that $\frac{4\lambda}{ma^2} > \frac{1 - q}{q}$, if h is sufficiently small, we get $C > 0$.

Let us define $y_0 = \frac{C}{B}$, $y_0 \in (0, 1)$ and if $y \leq y_0$, we obtain $\Phi_n(y) \geq 0$; if $y_0 \leq y \leq 1$, we obtain:

$$\Phi_n(y) \geq A\lambda^q y_0^{q+1} - B(1 - y_0)$$

.

This quantity will be positive if : $\lambda^q \geq \frac{B-C}{Ay_0^{q+1}}$.

Further, we have:

$$A \geq m(1 - \delta)^q \left(q + (1 - q)(1 - \delta) \left(1 - \frac{\Delta t_n}{T - t_n} \right) \right)$$

The solution at the time level t_{n+1} exists if $mq\Delta t_n \|\hat{u}_h^n\|_\infty < 1$, that is $\frac{\Delta t_n}{T - t_n} < \frac{1}{mq\lambda^q}$

and if $\lambda \geq \lambda_0 > \frac{1}{(mq)^{\frac{1}{q}}}$, we get: $A \geq mq(1 - \delta)^q$

and $\Phi_n(y)$ will be positive if:

$$(3.2) \quad \lambda^q \geq \frac{1}{mq(1-\delta)^q} \left(\frac{1}{q} + \frac{2\lambda}{a^2} + \frac{4\lambda}{ma^2}\delta \right) \left(\frac{1 + \frac{4\lambda}{ma^2} + \frac{2\lambda}{a^2}}{\frac{4\lambda}{ma^2}(1-\delta) - \delta \left(1 + \frac{2\lambda}{ma^2} \right) - \frac{1-q}{q}} \right)^{q+1}$$

The second member is a function of $\frac{\lambda}{a^2}$, hence if $\frac{\lambda}{a^2}$ is fixed such that $\frac{4\lambda}{ma^2} > \frac{1-q}{q}$, the inequality (3.2) will be satisfied if λ is large enough and h sufficiently small. Hence if the initial condition satisfies $u_{0h}(x) \geq \frac{\lambda}{T^{\frac{1}{q}}}\theta_h(\xi_0)$, λ and a satisfying (3.2), the solution blows up in finite time.

3.2. Blow-up of the solution.

Theorem 3.1. *Let $0 < q < 1$. The solution of problem (1.2) blows up in finite time.*

Proof: Since $u_0(x) \not\equiv 0$, there exists $\rho > 0$, $\epsilon > 0$, x_0 such that $u_0(x) \geq \epsilon > 0$ for x such that $|x - x_0| < \rho$. So we can choose T large enough such that $\frac{\lambda}{T^{\frac{1}{q}}} < \epsilon$ and $\frac{a}{T^{\frac{1-q}{2q}}} < \rho$.

We have $u_0(x) \geq \frac{\lambda}{T^{\frac{1}{q}}}\theta_h\left(\frac{x-x_0}{\zeta_0}\right)$ and then the solution blows up at time $T_0 \leq T^*$ where

$$T^* = \text{Max} \left(\left(\frac{\lambda}{\epsilon} \right)^q, \left(\frac{a}{\rho} \right)^{\frac{2q}{1-q}} \right).$$

In Fig1, we present the evolution of an initial condition u_0 for $m = 1$, $p = 1.5$. The solution blows up in finite time.




FIGURE 1. blow-up for $m=1$, $p=1,5$

REFERENCES

- [1] C. Bandle and H. Brunner. Blowup in diffusion equations: A survey. *J. Comput. Appl. Math.*, 97:3–22, 1998.
- [2] J. Bebernes and V.A. Galaktionov. On classification of blow-up patterns for a quasilinear heat equation. *Differential and Integral Equations*, 9(4):655–670, 1996.
- [3] V.A. Galaktionov. On localization conditions for unbounded solutions of quasilinear parabolic equations. *Sov.Math.Dokl.*, 1982.
- [4] V.A. Galaktionov. Best possible upper bounds for blowup solutions of the quasilinear heat conduction equation with source. *SIAM J. Math. Anal.*, 22(5):1293–1302, 1991.
- [5] V.A. Galaktionov. On a blow-up set for the quasilinear heat equation $u_t = (u^\sigma u_x)_x + u^{\sigma+1}$. *J. Differential Equations*, 101:66–79, 1993.
- [6] V.A. Galaktionov. Asymptotic behaviour of unbounded solutions of the nonlinear parabolic equation $u_t = (u^\sigma u_x)_x + u^\beta u$. *Differential Equations (English Translation)*, 21:751–758, 1985. *Differentsial'nye Uravneniya*, 21,n7,(1985) 1126–1134.
- [7] A.Y. Le Roux and M.N. Le Roux. Numerical solution of a nonlinear reaction diffusion equation. *Journal of Computational and Applied Mathematics*, 173:211–237, 2005.
- [8] A.Y. Le Roux and M.N. Le Roux. Numerical solution of a cauchy problem for nonlinear reaction diffusion processes. www-gm3.univ-mrs.fr, 2007.
- [9] P.E. Sacks. Global behavior for a class of nonlinear evolution equations. *SIAM J. Math. Anal.*, 16(2):233–250, 1985.
- [10] A.A. Samarskii, V.A. Galaktionov, S.P. Kurdyumov, and A.P. Mikhailov. *Blow-up in Quasilinear Parabolic Equations*. Walter de Gruyter, Berlin (English traduction), 1995. Nauka, Moscow(1987), in Russian.

IMB, INSTITUT DE MATHÉMATIQUES DE BORDEAUX, UMR5251, UNIVERSITÉ BORDEAUX1, 33405 TALENCE CEDEX, FRANCE

E-mail address: Marie-Noelle.Leroux@math.u-bordeaux1.fr